

If you have any questions or need assistance, please reach out to technical@bit.bio and we will do our best to support you.

Published June 2023, version 2

MATERIALS AND EQUIPMENT

- **ioMicroglia cells (cat no: io1021)**
Cultured to day 10 in a 96-well plate seeded at 37,000 cells/cm² (200µL/well)
- **comp:MM2x medium**
As described in the ioMicroglia user manual
- **comp:MM2x+LPS+IFN γ medium**
As described in Appendix 1
- **96-well U bottom plates**
- **Adhesive film**
- **LPS (50µg/mL)**
- **IFN γ (50µg/mL)**
- **Biological safety cabinet**
- **Standard tissue culture wares (pipettes, tips, culture plates)**
- **Normoxic cell culture incubator (37°C, 5% CO₂)**
- **Vortex**

ioMicroglia

Stimulation for cytokine release

Introduction

Microglia, as residents of the central nervous system, thrive in a complex and heterogeneous environment that is populated with neurons and glial cells. In their capacity as brain-resident immune cells, microglia play a critical role in responding to tissue damage and infection through the release of cytokines and phagocytosis of pathogens.


There is a need for functional and consistent human microglia for neuroimmune research and drug development for neurodegenerative diseases. Current models have constraints such as limitations in supply, complex protocols, or lot variability. All which raise challenges for drug discovery due to the generation of unreliable data and long experimental timelines.

ioMicroglia are human induced pluripotent stem cell (iPSC)-derived microglia, precision reprogrammed using opti-ox™ technology. At 10 days post-revival, they demonstrate maturity reminiscent of primary adult and foetal microglia. Functionally, ioMicroglia secrete proinflammatory and antiinflammatory cytokines (IL-6, TNF α , IL-1b, IL-12p70, IL-8, and IL-10) in response to stimuli such as Lipopolysaccharides (LPS) and Interferon Gamma (IFN γ). The protocol depicted below outlines general guidelines for stimulating cytokine release.

NOTES

- The following protocol recommends general guidelines. We encourage users to optimise the critical steps according to their experimental conditions.

Protocol overview



Protocol

This protocol is split into two sections:

1. Treatment of cells with LPS and IFN γ .
2. Harvesting of cell supernatants for cytokine readout.

1. Treatment of cells with LPS and IFN γ

- 1.1. Add comp:MM2x+LPS+IFN γ medium and comp:MM2x medium to separate reagent reservoirs.
- 1.2. Remove 100 μ L of expended culture medium from each well.
- 1.3. Add 100 μ L of the comp:MM2x+LPS+IFN γ medium to the treatment wells, for a final volume of 200 μ L.
- 1.4. Add 100 μ L of fresh, pre-warmed comp:MM2x medium to the control wells.
- 1.5. Incubate plates for 24 hours in an incubator at 37°C, 5% CO $_2$.

2. Harvesting of cell supernatants for cytokine readout

- 2.1. After the 24 hour incubation, remove plates from the incubator and transfer them to a biological safety cabinet.
- 2.2. Gently transfer 100 μ L of the medium from the culture plates to a fresh 96-well U bottom plate.
- 2.3. Cover with adhesive film and transfer to the -20°C freezer until ready to analyse on an Meso Scale Discovery (MSD) or ELISA based assay system.

Optional: The attached cells may be fixed for analysis via immunofluorescent staining at a later date.

Note: bit.bio uses an MSD system and examines the following cytokines: IL-6, TNF α , IL-1b, IL-12p70, IL-8, and IL-10.

Appendix 1

Preparation of comp:MM2x+LPS+IFN γ medium

1. Dilute 10 μ L LPS stock (50 μ g/mL) in 90 μ L of comp:MM2x medium.
2. Dilute 10 μ L IFN γ stock (50 μ g/mL) in 90 μ L of comp:MM2x medium.
3. Use the diluted reagents to prepare comp:MM2x+LPS+IFN γ medium as indicated in Table 1 (make 100 μ L per test well plus an excess ~10% for volume loss).
4. Vortex the comp:MM2x+LPS+IFN γ medium for 5 seconds.

Table 1- Preparation of comp:MM2x+LPS+IFN γ medium

Reagent	Stock concentration	Final concentration	comp:MM2x+LPS+IFN γ (for 1000 μ L)
comp:MM2x medium	-	-	976 μ L
Diluted LPS	5 μ g/mL	20ng/mL	4 μ L
Diluted IFN γ	5 μ g/mL	100ng/mL	20 μ L